

ZIBELINE INTERNATIONAL
PUBLISHINGISSN: 2521-0831 (Print)
ISSN: 2521-084X (Online)
CODEN: MSMADH

RESEARCH ARTICLE

ANALYTIC APPROXIMATE SOLUTION OF RABIES TRANSMISSION DYNAMICS USING HOMOTOPY PERTURBATION METHOD

Muhammad Sinan*

Department of Mathematics and Statistics, University of Swat
*Corresponding Author Email: sinanmathematics@gmail.com

This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

ARTICLE DETAILS

Article History:

Received 20 February 2020
Accepted 22 March 2020
Available online 09 April 2020

ABSTRACT

In this paper, we consider a mathematical model of Rabies disease which is an infectious disease. The model we are considering is a system of nonlinear ordinary differential equations and it is difficult to find an exact solution. He's Homotopy perturbation method is employed to compute an approximation to the solution of the system of nonlinear ordinary differential equations. The findings obtained by HPM are compared with a nonstandard finite difference (NSFD) and Runge-Kutta fourth order (RK4) methods. Some plots are presented to show the reliability and simplicity of the method.

KEYWORDS

Mathematical model, infectious, nonlinear ordinary, homotopy, equations.

1. INTRODUCTION

Rabies is an infection that mostly affects the brain of an Infected animal or individual, caused by viruses belonging to The genus *Lyssavirus* of the family *Rhabdoviridae* and order *Mononegavirales* (Hayman et al., 2011). This disease has become a global threat and it is also estimated that rabies occurs in more than 150 countries and territories (Rabies fact sheet, 2016). Raccoons, skunks, bats, and foxes are the main animals that transmit the virus in the United States. In Asia, Africa, and Latin America, it is known that dogs are the main source of transmission of the rabies virus into the human population (Zhang et al., 2011; Grace, 2014; Wiraningih et al., 2015; Global Alliance for rabies control, 2016). When the rabies virus enters the human body or that of an animal, the infection (virus) moves rapidly along the neural pathways to the central nervous system; from there the virus continues to spread to other organs and causes injury by interrupting various nerves (Rabies fact sheet, 2016; CIA, 2016; Birkho and Rota, 1962). The symptoms of rabies are quite similar to those of encephalitis (Rupprecht et al., 2010). Due to the movement of dogs in homes or the surroundings, the risk of being infected by a rabid dog can never be guaranteed. Rabies is a major health problem in many populations dense with dogs, especially in areas where there are less or no preventive measures (vaccination and treatment) for dogs and humans (Lakshmikantham et al., 1994; Diekmann et al., 1990; Pielou, 1969; May, 1973; Martcheva, 2015). Treatment after exposure to the rabies virus is known as post-exposure Prophylaxis (PEP) and vaccination before exposure to the infection is known as pre-exposure prophylaxis.

Since most of the mathematical models raised from biological problems are nonlinear by nature and it is difficult to find the analytical solution of such problems (Yusuf and Benyah, 2012; Hove-Musekwa et al., 2011; Deleon, 2009; Lasalle, 1976). Therefore, it is a great challenge for mathematicians and researchers to find such numerical and perturbation

methods which give the best approximation to the solution of such nonlinear problems. Convergence and accuracy are the key concepts while developing and implementing a numerical scheme otherwise results will be inappropriate. As far as the analytical perturbation methods are concerned, a parameter (negligibly small) needs to be exerted in the equation. Exertion and production of such parameter is a difficult task in these methods.

Recent research provided powerful methods like artificial parameter method in which this small parameter is absent. An approximate solution of nonlinear differential equations can be effectively obtained using the well-known Homotopy Analysis Method (HAM) (Zhang et al., 2015; Sharomi and Malik, 2017; Ali et al., 2018; Bushnaq et al., 2018a; Bushnaq et al., 2017; Bushnaq et al., 2018b). The method is used with perturbation methods in recent decades. The basic and fundamental scheme of the method was first introduced by Liao and He. The method uses a free parameter whose appropriate selection yields fast convergence of the algorithm. At the initial stage, He introduced HPM and applies the procedure to some interesting problems (Rupprecht et al., 2010; Aubert, 1999; Shah et al., 2018; Asamoah et al., 2017; Samia et al., 2018). Another group researchers used optimal Homotopy Analysis Method (OHAM) in order to obtain the solution of multi-point boundary value problem (Ali et al., 2017).

The methods mentioned above are free from the choice of small parameter and have all the advantages of perturbation methods. This work is an extension by considering HPM applied to leptospirosis epidemic model (Aubert, 1999; Anderson and May, 1982; Bohrer et al., 2002; Levin et al., 2012; Coyne et al., 1989; Childs et al., 2000; Hampson et al., 2007; Carroll et al., 2010). We will compare the results obtained by HPM with Runge-Kutta fourth order (RK4) method. Numerous problems of nonlinear nature can be solved accurately and effectively using HPM because of its rapid convergence (Wang and Lou, 2008; Yang and Lou, 2009; Zinsstag et

Quick Response Code



Access this article online

Website:
www.matrixscience.comDOI:
10.26480/msmk.01.2020.01.05

al., 2009; Ding et al., 2007; Smith and Cheeseman, 2002; Tchaenche and Bauch, 2012). The rest of the manuscript is as follow. In Section 2, we included the basic concept of HPM. In Section 3, the model is formulated and solved by HPM. In section 4, some numerical result and discussion is given, and the conclusion is presented at the end of the paper.

2. ANALYSIS OF HOMOTOPY PERTURBATION METHOD (HPM)

To illustrate the basic idea of HPM, consider the general nonlinear differential equation

$$A(\mu) - f(r) = 0, r \in \Omega, \tag{1}$$

with the boundary condition,

$$\beta \left(\mu, \frac{\partial \mu}{\partial n} \right) = 0, r \in \Gamma, \tag{2}$$

where A is a general differential operator, β is a boundary operator, $f(r)$ a known analytic function, Γ is the boundary of the domain Ω . The operator A is divided into linear part L and nonlinear part N . Therefore, equation (1) can be written as,

$$L(u) + N(u) - f(r) = 0 \tag{3}$$

By using the Homotopy technique, one can construct a Homotopy

$$v(r, p): \Omega \times [0,1] \rightarrow R \tag{4}$$

which satisfies

$$\begin{aligned} H(v, p) &= (1 - p)[L(v) - L(\mu_0)] \\ + p[A(v - f(r))] &= 0 \end{aligned} \tag{5}$$

or

$$\begin{aligned} H(v, p) &= L(v) - L(\mu_0) \\ + pL(v_0) + p[N(v) - f(r)] &= 0 \end{aligned} \tag{6}$$

Where, $p \in [0; 1]$ is an embedding parameter and μ_0 is the initial approximation of the given equation that satisfies the boundary conditions. Clearly, we have

$$H(v, 0) = L(v) - L(\mu_0) = 0 \tag{7}$$

$$H(v, 1) = A(v) - f(r) = 0 \tag{8}$$

The changing process of p from zero to one is just that of $v(r, p)$ changing from $\mu_0(r)$ to $\mu(r)$. This is called deformation, and $L(v) - L(\mu_0)$ and $A(v) - f(r)$ are called homotopic in topology. If the embedding parameter $p(0 \leq p \leq 1)$ is considered as a small parameter, applying the classical perturbation technique, we can naturally assume that the solution of the equation can be given as a power series in p ,

$$v = v_0 + pv_1 + p^2v_2 + p^3v_3 + \dots \tag{9}$$

Setting $p = 1$ results in the approximate solution as

$$v = \lim_{p \rightarrow 1} v = v_0 + v_1 + v_2 + v_3 + \dots \tag{10}$$

3. THE MODEL OF ORDINARY DIFFERENTIAL EQUATIONS

$$\begin{aligned} \frac{dS_D}{dt} &= A_D - (1 - v_D)\beta_{DD}S_DI_D - (m_D + v_D)S_D + \delta\varepsilon_DE_D + \alpha_DR_D, \\ \frac{dE_D}{dt} &= (1 - v_D)\beta_{DD}S_DI_D - ((1 - \rho_D)\delta\gamma_D + m_D + \rho_D + \delta\varepsilon_D + C_D)E_D, \\ \frac{dI_D}{dt} &= (1 - \rho_D)\delta\gamma_DE_D - (m_D + \mu_D)I_D, \\ \frac{dR_D}{dt} &= v_DS_D + \rho_DE_D - (m_D + \alpha_D)R_D, \\ \frac{dS_H}{dt} &= B_H - (1 - v_H)\beta_{DH}S_HI_D - (m_H + v_H)S_H + \delta_H\varepsilon_HE_H + \alpha_HR_H, \\ \frac{dE_H}{dt} &= (1 - v_H)\beta_{DH}S_HI_D - ((1 - \rho_H)\delta_H\gamma_H + m_H + \rho_H + \delta_H\varepsilon_H)E_H, \\ \frac{dI_H}{dt} &= (1 - \rho_H)\delta_H\gamma_HE_H - (m_H + \mu_H)I_H, \end{aligned} \tag{11}$$

$$\frac{dR_H}{dt} = v_HS_H + \rho_HE_H - (m_H + \alpha_H)R_H.$$

With

$$\begin{aligned} S_D(0) > 0, E_D(0) \geq 0, I_D(0) \geq 0, R_D(0) \geq 0, \\ S_H(0) > 0, E_H(0) \geq 0, I_H(0) > 0, R_H(0) > 0. \end{aligned} \tag{12}$$

To find out the solution of the given model by Homotopy Perturbation Method (HPM). Let us consider the Homotopy which becomes for the system (1), such that,

$$\begin{aligned} LS_D(t) - LS_{D0}(t) &= p[A_D - (1 - v_D)\beta_{DD}S_DI_D - (m_D + v_D)S_D + \delta\varepsilon_DE_D + \alpha_DR_D - LS_{D0}(t)] \\ LE_D(t) - LE_{D0}(t) &= p[(1 - v_D)\beta_{DD}S_DI_D - (1 - \rho_D)\delta\gamma_D + m_D + \rho_D + \delta\varepsilon_D + C_D)E_D - LE_{D0}(t)] \\ LI_D(t) - LI_{D0}(t) &= p[(1 - \rho_D)\delta\gamma_DE_D - (m_D + \mu_D)I_D - LI_{D0}(t)] \\ LR_D(t) - LR_{D0}(t) &= p[v_DS_D + \rho_DE_D - (m_D + \alpha_D)R_D - LR_{D0}(t)] \\ LS_H(t) - LS_{H0}(t) &= p[B_H - (1 - v_H)\beta_{DH}S_HI_D - (m_H + v_H)S_H + \delta_H\varepsilon_HE_H + \alpha_HR_H - LS_{H0}(t)] \\ LE_H(t) - LE_{H0}(t) &= p[(1 - v_H)\beta_{DH}S_HI_D - ((1 - \rho_H)\delta_H\gamma_H + m_H + \rho_H + \delta_H\varepsilon_H)E_H - LE_{H0}(t)] \\ LI_H(t) - LI_{H0}(t) &= p[\delta_H\gamma_HE_H - (m_H + \mu_H)I_H - LI_{H0}(t)] \\ LR_H(t) - LR_{H0}(t) &= p[v_HS_H + \rho_HE_H - (m_H + \alpha_H)R_H - LR_{H0}(t)] \end{aligned}$$

Assume that the solution of model (1) is in the form,

$$\begin{aligned} S_D(t) &= S_{D0} + pS_{D1} + p^2S_{D2} + \dots \\ E_D(t) &= E_{D0} + pE_{D1} + p^2E_{D2} + \dots \\ I_D(t) &= I_{D0} + pI_{D1} + p^2I_{D2} + \dots \\ R_D(t) &= R_{D0} + pR_{D1} + p^2R_{D2} + \dots \\ S_H(t) &= S_{H0} + pS_{H1} + p^2S_{H2} + \dots \\ E_H(t) &= E_{H0} + pE_{H1} + p^2E_{H2} + \dots \\ I_H(t) &= I_{H0} + pI_{H1} + p^2I_{H2} + \dots \\ R_H(t) &= R_{H0} + pR_{H1} + p^2R_{H2} + \dots \end{aligned} \tag{13}$$

Now using $S_D(t)$, $E_D(t)$, $I_D(t)$, $R_D(t)$, $S_H(t)$, $E_H(t)$, $I_H(t)$, and $R_H(t)$ in equation (3) and compare the coefficients of zeroth order p^0 , first order p^1 and second order p^2 perturbation and so on respectively in each equation of system (13), Comparing the coefficient of p^0, p^1, p^2, \dots , such that,

$$\begin{aligned} p^1: LS_{D1} &= A_D - (1 - v_D)\beta_{DD}S_{D0}I_{D0} - (m_D - v_D)S_{D0} + \delta\varepsilon_DR_{D0} + \alpha_DR_{D0}, \\ p^1: LE_{D1} &= (1 - v_D)\beta_{DD}S_{D0}I_{D0} - ((1 - \rho_D)\delta\gamma_D + m_D + \rho_D + \delta\varepsilon_D + C_D)E_{D0}, \\ p^1: LI_{D1} &= (1 - \rho_D)\delta\gamma_DE_{D0} - (m_D + \mu_D)I_{D0}, \\ p^1: LR_{D1} &= v_DS_{D0} - (m_D + \mu_D)I_{D0}, \\ p^1: LS_{H1} &= B_H - (1 - \gamma_H)\beta_{DH}\delta_H I_{D0} - (m_H + \gamma_H)S_{H0} + \beta_H + \varepsilon_H E_{H0} + \alpha_H R_{H0}, \\ p^1: LE_{H1} &= (1 - \rho_H)\beta_{DH}S_{H0}I_{D0} - ((1 - \rho_H)(\delta_H\gamma_H + m_H + \rho_H + \delta_H\varepsilon_D))E_{H0}, \\ p^1: LI_{H1} &= (1 - \rho_H)\delta_H\gamma_HE_{H0} - (m_H + \mu_H)I_{H0}, \\ p^1: LR_{H1} &= \rho_HS_{H0} + \rho_HE_{H0} - (m_H + \alpha_H)R_{H0}, \end{aligned} \tag{14}$$

and

$$\begin{aligned} p^2: LS_{D2} &= -(1 - v_D)\beta_{DD}S_{D1}I_{D1} - (m_D + v_D)S_{D1} + \delta\varepsilon_DR_{D1}, \\ p^2: LE_{D2} &= (1 - v_D)\beta_{DD}S_{D1}I_{D1} - ((1 - \rho_D)\delta\gamma_D + m_D + \rho_D + \delta\varepsilon_D + C_D), \\ p^2: LI_{D2} &= (1 - \rho_D)\delta\gamma_DE_{D1} - (m_D + \mu_D)I_{D1}, \\ p^2: LR_{D2} &= v_DS_{D1} - E_{D1} - (m_D + \alpha_D)R_{D1}, \\ p^2: LS_{H2} &= -(1 - \gamma_H)\beta_{DH}S_{H1}I_{D1} - (m_H + v_H)S_{H1} + \varepsilon_HE_{H1} + \alpha_HR_{H1}, \\ p^2: LE_{H2} &= (1 - \rho_H)\beta_{DH}S_{H1}I_{D1} - ((1 - \rho_H)(\delta_H\gamma_H + m_H + \rho_H + \delta_H\varepsilon_H)E_{H1}), \\ p^2: LI_{H2} &= (1 - \rho_H)\delta_H\gamma_HE_{H1} - (m_H + \mu_H)I_{H1}, \\ p^2: LR_{H2} &= v_HS_{H1} + \rho_HE_{H1} - (m_H + \alpha_H)R_{H1} \end{aligned} \tag{15}$$

In order to obtain the solution of the problem, we consider the following cases,

3.1 Zeroth Order Problem or P^0

$$\begin{aligned} S_{D0}(t) &= 44, E_{D0}(t) = 88, \\ I_{D0}(t) &= 100, R_{D0}(t) = 110 \\ S_{H0}(t) &= 105, E_{H0}(t) = 103, \end{aligned} \tag{16}$$

$$I_{H0}(t) = 90, R_{H0}(t) = 95$$

3.2 First Order Problem or P¹

$$\begin{aligned} S_{D1}(t) &= (A_D - (1 - v_D)\beta_{D0}S_{D0}I_{D0} - (m_D - v_D)S_{D0} + \delta\varepsilon_D R_{D0} + \alpha_D R_{D0})t, \\ E_{D1}(t) &= ((1 - v_D)\beta_{D0}S_{D0} - ((1 - \rho_D)\delta\gamma_D + m_D + \rho_D + \delta\varepsilon_D + C_D)E_{D0})t, \\ I_{D1}(t) &= ((1 - \rho_D)\delta\gamma_D E_{D0} - (m_D + \mu_D)I_{D0})t, \\ R_{D1}(t) &= (v_D S_{D0} - (m_D + \mu_D)I_{D0})t, \\ S_{H1}(t) &= (B_H - (1 - \gamma_H)\beta_{DH}\delta_H I_{D0} - (m_H + \gamma_H)S_{H0} + \beta_H + \varepsilon_H E_{H0} + \alpha_H R_{H0})t, \\ E_{H1}(t) &= ((1 - \rho_H)\beta_{DH}S_{H0}I_{D0} - ((1 - \rho_H)(\delta_H\gamma_H + m_H + \rho_H + \delta_H\varepsilon_D))E_{H0})t, \\ I_{H1}(t) &= ((1 - \rho_H)\delta_H\gamma_H E_{H0} - (m_H + \mu_H)I_{H0})t, \\ R_{H1}(t) &= (\rho_H S_{H0} + \rho_H E_{H0} - (m_H + \alpha_H)R_{H0})t, \end{aligned} \tag{17}$$

3.3 Second Order Problem or P²

$$\begin{aligned} S_{D2}(t) &= -(1 - u_D)\beta_{DD}((A_D - (1 - v_D)\beta_{D0}S_{D0}I_{D0} - (m_D - v_D)S_{D0} + \delta\varepsilon_D R_{D0} + \alpha_D R_{D0})\frac{t^2}{2} \\ E_{D2}(t) &= (1 - v_D)\beta_{DD}((A_D - (1 - v_D)\beta_{D0}S_{D0}I_{D0} - (m_D - v_D)S_{D0} + \delta\varepsilon_D R_{D0} + \alpha_D R_{D0})\frac{t^2}{2} \\ &\quad + ((1 - \rho_D)\delta\gamma_D E_{D0} - (m_D + \mu_D)I_{D0})\frac{t^2}{2} - ((1 - \rho_D)\delta\gamma_D + m_D + \rho_D + \delta\varepsilon_D + C_D) \\ I_{D2}(t) &= (1 - \rho_D)\delta\gamma_D ((1 - v_D)\beta_{DD}S_{D0} - ((1 - \rho_D)\delta\gamma_D + m_D + \rho_D + \delta\varepsilon_D + C_D)E_{D0})\frac{t^2}{2} - (m_D + \mu_D)((1 - \rho_D)\delta\gamma_D E_{D0} - (m_D + \mu_D)I_{D0})\frac{t^2}{2} \\ R_{D2}(t) &= v_D(A_D - (1 - v_D)\beta_{D0}S_{D0}I_{D0} - (m_D - v_D)S_{D0} + \delta\varepsilon_D R_{D0} + \alpha_D R_{D0})\frac{t^2}{2} - ((1 - v_D)\beta_{DD}S_{D0} - ((1 - \rho_D)\delta\gamma_D + m_D + \rho_D + \delta\varepsilon_D + C_D)E_{D0})\frac{t^2}{2} - (m_D + \alpha_D)(v_D S_{D0} - (m_D + \mu_D)I_{D0})\frac{t^2}{2} \\ S_{H2}(t) &= -(1 - \gamma_H)\beta_{DH}((B_H - (1 - \gamma_H)\beta_{DH}\delta_H I_{D0} - (m_H + \gamma_H)S_{H0} + \beta_H + \varepsilon_H E_{H0} + \alpha_H R_{H0})\frac{t^2}{2} - ((1 - \rho_H)\delta_H\gamma_H E_{H0} - (m_H + \mu_H)I_{H0})\frac{t^2}{2} - (m_H + v_H)((1 - \rho_H)\beta_{DH}S_{H0}I_{D0} - ((1 - \rho_H)(\delta_H\gamma_H + m_H + \rho_H + \delta_H\varepsilon_D))E_{H0})\frac{t^2}{2} + \varepsilon_H((1 - \rho_H)\beta_{DH}S_{H0}I_{D0} - ((1 - \rho_H)(\delta_H\gamma_H + m_H + \rho_H + \delta_H\varepsilon_D))E_{H0})\frac{t^2}{2} + \alpha_H(\rho_H S_{H0} + \rho_H E_{H0} - (m_H + \alpha_H)R_{H0})\frac{t^2}{2} \\ E_{H2}(t) &= (1 - \rho_H)\beta_{DH}((B_H - (1 - \gamma_H)\beta_{DH}\delta_H I_{D0} - (m_H + \gamma_H)S_{H0} + \beta_H + \varepsilon_H E_{H0} + \alpha_H R_{H0})\frac{t^2}{2} - ((1 - \rho_H)\delta_H\gamma_H E_{H0} - (m_H + \mu_H)I_{H0})\frac{t^2}{2} - ((1 - \rho_H)(\delta_H\gamma_H + m_H + \rho_H + \delta_H\varepsilon_D))E_{H0})\frac{t^2}{2} \\ I_{H2}(t) &= (1 - \rho_H)\delta_H\gamma_H ((1 - \rho_H)\beta_{DH}S_{H0}I_{D0} - ((1 - \rho_H)(\delta_H\gamma_H + m_H + \rho_H + \delta_H\varepsilon_D))E_{H0})\frac{t^2}{2} - (m_H + \mu_H)((1 - \rho_H)\delta_H\gamma_H E_{H0} - (m_H + \mu_H)I_{H0})\frac{t^2}{2} \\ R_{H2}(t) &= v_H(B_H - (1 - \gamma_H)\beta_{DH}\delta_H I_{D0} - (m_H + \gamma_H)S_{H0} + \beta_H + \varepsilon_H E_{H0} + \alpha_H R_{H0})\frac{t^2}{2} + \rho_H((1 - \rho_H)\beta_{DH}S_{H0}I_{D0} - ((1 - \rho_H)(\delta_H\gamma_H + m_H + \rho_H + \delta_H\varepsilon_D))E_{H0})\frac{t^2}{2} - (m_H + \alpha_H)(\rho_H S_{H0} + \rho_H E_{H0} - (m_H + \alpha_H)R_{H0})\frac{t^2}{2} \end{aligned} \tag{18-25}$$

To find the solution we consider $p = 1$ in the system (14), we get

$$\begin{aligned} S_D(t) &= S_{D0} + S_{D1} + S_{D2} + \dots \\ E_D(t) &= E_{D0} + E_{D1} + E_{D2} + \dots \\ I_D(t) &= I_{D0} + I_{D1} + I_{D2} + \dots \\ R_D(t) &= R_{D0} + R_{D1} + R_{D2} + \dots \\ S_H(t) &= S_{H0} + S_{H1} + S_{H2} + \dots \\ E_H(t) &= E_{H0} + E_{H1} + E_{H2} + \dots \\ I_H(t) &= I_{H0} + I_{H1} + I_{H2} + \dots \\ R_H(t) &= R_{H0} + R_{H1} + R_{H2} + \dots \end{aligned} \tag{26}$$

Thus, we have a solution to the model below, i.e

$$S_D(t) = 44 + (A_D - (1 - v_D)\beta_{D0}S_{D0}I_{D0} - (m_D - v_D)S_{D0} + \delta\varepsilon_D R_{D0} + \alpha_D R_{D0})t + ((1 - u_D)\beta_{DD}((A_D - (1 - v_D)\beta_{D0}S_{D0}I_{D0} - (m_D - v_D)S_{D0} + \delta\varepsilon_D R_{D0} + \alpha_D R_{D0})\frac{t^2}{2} - ((1 - \rho_D)\delta\gamma_D + m_D + \rho_D + \delta\varepsilon_D + C_D)E_{D0})\frac{t^2}{2} + \dots$$

$$\begin{aligned} &\delta\varepsilon_D R_{D0} + \alpha_D R_{D0})\frac{t^2}{2} - ((1 - \rho_D)\delta\gamma_D E_{D0} - (m_D + \mu_D)I_{D0})\frac{t^2}{2} - \\ &(m_D + v_D)(A_D - (1 - v_D)\beta_{D0}S_{D0}I_{D0} - (m_D - v_D)S_{D0} + \delta\varepsilon_D R_{D0} + \\ &\alpha_D R_{D0})\frac{t^2}{2} + \delta(v_D S_{D0} - (m_D + \mu_D)I_{D0})\frac{t^2}{2} + \dots \end{aligned} \tag{27}$$

$$\begin{aligned} E_D(t) &= 88 + ((1 - v_D)\beta_{DD}S_{D0} - ((1 - \rho_D)\delta\gamma_D + m_D + \rho_D + \delta\varepsilon_D + C_D)E_{D0})t + ((1 - v_D)\beta_{DD}((A_D - (1 - v_D)\beta_{D0}S_{D0}I_{D0} - (m_D - v_D)S_{D0} + \delta\varepsilon_D R_{D0} + \alpha_D R_{D0})\frac{t^2}{2} - ((1 - \rho_D)\delta\gamma_D + m_D + \rho_D + \delta\varepsilon_D + C_D)E_{D0})\frac{t^2}{2} + \dots \end{aligned} \tag{28}$$

$$\begin{aligned} I_D(t) &= 100 + ((1 - \rho_D)\delta\gamma_D E_{D0} - (m_D + \mu_D)I_{D0})t + ((1 - \rho_D)\delta\gamma_D ((1 - v_D)\beta_{DD}S_{D0} - ((1 - \rho_D)\delta\gamma_D + m_D + \rho_D + \delta\varepsilon_D + C_D)E_{D0})\frac{t^2}{2} - \\ &(m_D + \mu_D)((1 - \rho_D)\delta\gamma_D E_{D0} - (m_D + \mu_D)I_{D0})\frac{t^2}{2} + \dots \end{aligned} \tag{29}$$

$$\begin{aligned} R_D(t) &= 110 + (v_D S_{D0} - (m_D + \mu_D)I_{D0})t + (v_D(A_D - (1 - v_D)\beta_{D0}S_{D0}I_{D0} - (m_D - v_D)S_{D0} + \delta\varepsilon_D R_{D0} + \alpha_D R_{D0})\frac{t^2}{2} - ((1 - v_D)\beta_{DD}S_{D0} - \\ &((1 - \rho_D)\delta\gamma_D + m_D + \rho_D + \delta\varepsilon_D + C_D)E_{D0})\frac{t^2}{2} - (m_D + \alpha_D)(v_D S_{D0} - \\ &(m_D + \mu_D)I_{D0})\frac{t^2}{2} + \dots \end{aligned} \tag{30}$$

$$\begin{aligned} S_H(t) &= 105 + (B_H - (1 - \gamma_H)\beta_{DH}\delta_H I_{D0} - (m_H + \gamma_H)S_{H0} + \beta_H + \varepsilon_H E_{H0} + \alpha_H R_{H0})t + ((1 - \gamma_H)\beta_{DH}((B_H - (1 - \gamma_H)\beta_{DH}\delta_H I_{D0} - (m_H + \gamma_H)S_{H0} + \beta_H + \varepsilon_H E_{H0} + \alpha_H R_{H0})\frac{t^2}{2} - ((1 - \rho_H)\delta_H\gamma_H E_{H0} - (m_H + \mu_H)I_{H0})\frac{t^2}{2} - (m_H + v_H)((1 - \rho_H)\beta_{DH}S_{H0}I_{D0} - ((1 - \rho_H)(\delta_H\gamma_H + m_H + \rho_H + \delta_H\varepsilon_D))E_{H0})\frac{t^2}{2} + \varepsilon_H((1 - \rho_H)\beta_{DH}S_{H0}I_{D0} - ((1 - \rho_H)(\delta_H\gamma_H + m_H + \rho_H + \delta_H\varepsilon_D))E_{H0})\frac{t^2}{2} + \alpha_H(\rho_H S_{H0} + \rho_H E_{H0} - (m_H + \alpha_H)R_{H0})\frac{t^2}{2} + \dots \end{aligned} \tag{31}$$

$$\begin{aligned} E_H(t) &= 103 + ((1 - \rho_H)\beta_{DH}S_{H0}I_{D0} - ((1 - \rho_H)(\delta_H\gamma_H + m_H + \rho_H + \delta_H\varepsilon_D))E_{H0})t + ((1 - \rho_H)\beta_{DH}((B_H - (1 - \gamma_H)\beta_{DH}\delta_H I_{D0} - (m_H + \gamma_H)S_{H0} + \beta_H + \varepsilon_H E_{H0} + \alpha_H R_{H0})\frac{t^2}{2} - ((1 - \rho_H)\delta_H\gamma_H E_{H0} - (m_H + \mu_H)I_{H0})\frac{t^2}{2} - (m_H + v_H)((1 - \rho_H)\beta_{DH}S_{H0}I_{D0} - ((1 - \rho_H)(\delta_H\gamma_H + m_H + \rho_H + \delta_H\varepsilon_D))E_{H0})\frac{t^2}{2} + \varepsilon_H((1 - \rho_H)\beta_{DH}S_{H0}I_{D0} - ((1 - \rho_H)(\delta_H\gamma_H + m_H + \rho_H + \delta_H\varepsilon_D))E_{H0})\frac{t^2}{2} + \alpha_H(\rho_H S_{H0} + \rho_H E_{H0} - (m_H + \alpha_H)R_{H0})\frac{t^2}{2} + \dots \end{aligned} \tag{32}$$


$$\begin{aligned} I_H(t) &= 90 + ((1 - \rho_H)\delta_H\gamma_H E_{H0} - (m_H + \mu_H)I_{H0})t + ((1 - \rho_H)\delta_H\gamma_H ((1 - \rho_H)\beta_{DH}S_{H0}I_{D0} - ((1 - \rho_H)(\delta_H\gamma_H + m_H + \rho_H + \delta_H\varepsilon_D))E_{H0})\frac{t^2}{2} - (m_H + \mu_H)((1 - \rho_H)\delta_H\gamma_H E_{H0} - (m_H + \mu_H)I_{H0})\frac{t^2}{2} + \dots \end{aligned} \tag{33}$$

$$\begin{aligned} R_H(t) &= 95 + (\rho_H S_{H0} + \rho_H E_{H0} - (m_H + \alpha_H)R_{H0})t + (v_H(B_H - (1 - \gamma_H)\beta_{DH}\delta_H I_{D0} - (m_H + \gamma_H)S_{H0} + \beta_H + \varepsilon_H E_{H0} + \alpha_H R_{H0})\frac{t^2}{2} + \rho_H((1 - \rho_H)\beta_{DH}S_{H0}I_{D0} - ((1 - \rho_H)(\delta_H\gamma_H + m_H + \rho_H + \delta_H\varepsilon_D))E_{H0})\frac{t^2}{2} - (m_H + \alpha_H)(\rho_H S_{H0} + \rho_H E_{H0} - (m_H + \alpha_H)R_{H0})\frac{t^2}{2} + \dots \end{aligned} \tag{34}$$

4. NUMERICAL RESULTS AND DISCUSSION

Table 1: Description of Parameter and its Value.		
Notation	Description of Parameters	Values
A_D	Rate at which S _D (t) is increased by recruitment	1.35
v_D	The control strategy due to public education and vaccination in the dogs	2.50
β_{DD}	Contact rate of the rabid dog into the dog	3.000
v_H	The preexposure prophylaxis(vaccination)	4.008
ρ_D	The rate at which exposed dogs are treated by their owners	9.020
C_D	The rate at which exposed dogs die due to culling	12.223
μ_D	The death rate associated with rabies infection in dogs	13.322
B_H	Birth or immigration rate into the susceptible human population	14.444

β_{DH}	Contact rate of infectious dogs to the human population	16.666
α_H	The rate of losing immunity in the human population	18.888
α_D	The rate of losing immunity in dogs' population	17.777
m_D	The natural death rate of dogs	19.999
m_H	The mortality rate of humans	20.222
ρ_H	The rate at which administrating treatment to affected humans	15.555
μ_H	The disease induces death in humans	23.333




5. CONCLUSION

In this article, we used a semi-analytical approach that is Homotopy Perturbation Method (HPM), compared with other kind of methods like Runge-Kutta fourth order (RK4) and NSFDF methods for the solution of optimal control model of rabies transmission dynamics in dogs and the best way of reducing death rate of rabies in humans. We concluded that the Homotopy perturbation method is a powerful technique to solve nonlinear differential equations.

ACKNOWLEDGEMENT

I would like to express my sincere gratitude to Dr. Amir Khan, Assistant Professor in Mathematics, University of Swat for the continuous support for solving this problem and study the related research, for his patience, motivation, and immense knowledge. His guidance helped me in all the time of research and writing of his paper. Once again thanks to Dr. Amir Khan.

ORCID iD

Muhammad Sinan  <https://orcid.org/0000-0003-2177-3806>

REFERENCES

- Ali, A., Laila, H., Shah, K., 2018. Analytical solution of General Fisher's Equation by using Laplace Adomian decomposition method, *J. Pur. Appl. Math.*, 2 (3), 1-4.
- Ali, A., Rabie, F., Shah, K., 2017. On Ulams type stability for a class of impulsive fractional differential equations with nonlinear integral boundary conditions. *Journal of Nonlinear Science and Applications*, 10, 4760-4775.
- Anderson, R.M., May, R.M., 1982. *Population biology of infectious diseases*. Proceedings of the Dahlem Workshop, Springer, Berlin, Germany.
- Asamoah, J.K.K., Oduro, F.T., Bonyah, E., Seidu, B., 2017. Modelling of Rabies Transmission Dynamics Using Optimal Control Analysis. *Journal*

of Applied Mathematics, Article ID 2451237, 23 page, 2017. <https://doi.org/10.1155/2017/2451237>.

- Aubert, M., 1999. Costs and benefits of rabies control in wildlife in France. *Revue Scientifique et Technique de IOIE*, 18 (2), Pp. 533-543.
- Birkho, G., Rota, G., 1962. *Ordinary differential equations*. Ginn and Co, New York, NY, USA.
- Bohrer, G., Shem-Tov, S., Summer, E., Or, K., Saltz, D., 2002. The effectiveness of various rabies spatial vaccination patterns in a simulated host population with clumped distribution. *Ecological Modelling*, 152 (2-3), Pp. 205-211.
- Bushnaq, S., Ali, S., Shah, K., Arif, M., 2018. Exact solution to nonlinear biological population model with fractional order. *Thermal Science Journal, Special Issue*, 10 pages.
- Bushnaq, S., Khan, S.A., Shah, K., Zaman, G., 2018. Existence theory of HIV-1 infection model by using arbitrary order derivative of without singular kernel type. *Journal of Mathematical Analysis*, 10 (9), Pp. 1-13.
- Bushnaq, S., Khan, S.A., Shah, K., Zaman, G., Liu, F., 2018. Mathematical analysis of HIV/AIDS infection model with Caputo-Fabrizio fractional derivative. *Cogent Mathematics & Statistics*, 5, Pp. 1432-1521.
- Carroll, M.J., Singer, A., Smith, G.C., Cowan, D.P., Massei, G., 2010. The use of immunosuppression to improve rabies eradication in urban dog populations. *Wildlife Research*, 37 (8), Pp. 676-687.
- Childs, J.E., Curns, A.T., Dey, M.E., 2000. Predicting the local dynamics of epizootic rabies among raccoons in the United States. *Proceedings of the National Academy of Sciences of the United States of America*, 97 (25), Pp. 13666-13671.
- CIA. 2016. *World factbook and other sources*.
- Coyne, M.J., Smith, G., McAllister, F.E., 1989. Mathematic model for the population biology of rabies in raccoons in the mid- Atlantic states. *American journal of veterinary research*, 50 (12), Pp. 2148-2154.
- De Leon, C.V., 2009. Constructions of Lyapunov functions for classics SIS, SIR and SIRS epidemic model with variable population size. *Foro-Red-Mat: Revista Electronica De Contenido Matematico*, 26 (5), article 1.
- Diekmann, O., Heesterbeek, J.A., Metz, J.A., 1990. On the definition and the computation of the basic reproduction ratio R_0 in models for infectious diseases in heterogeneous populations. *Journal of Mathematical Biology*, 28 (4), Pp. 365-382.
- Ding, W., Gross, L.J., Langston, K., Lenhart, S., Real, L.A., 2007. Rabies in raccoons: optimal control for a discrete time model on a spatial grid. *Journal of Biological Dynamics*, 1 (4), Pp. 379-393.
- Global alliance for rabies control. 2016. <https://rabiesalliance.org/>.
- Grace, O.A., 2014. Modelling of the spread of rabies with pre-exposure vaccination of humans. *Mathematical Theory and Modeling*, 4 (8).
- Hampson, K., Dusho, J., Bingham, J., Bruckner, G., Ali, Y.H., Dobson, A., 2007. Synchronous cycles of domestic dog rabies in sub-Saharan Africa and the impact of control efforts. *Proceedings of the National Academy of Sciences of the United States of America*, 104 (18), Pp. 7717-7722.
- Hayman, D.T.S., Johnson, N., Horton, D.L., 2011. Evolutionary history of rabies in Ghana. *PLoS Neglected Tropical Diseases*, 5 (4), Article ID e1001.
- Hove-Musekwa, S.D., Nyabadza, F., Mambili-Mamboundou, H., 2011. Modelling hospitalization, home-based care, and individual withdrawal for people living with HIV/AIDS in high prevalence settings. *Bulletin of Mathematical Biology*, 73 (12), Pp. 2888-2915.
- Lakshmikantham, V., Leela, S., Kaul, S., 1994. Comparison principle for impulsive differential equations with variable times and stability theory, *Nonlinear Analysis. Theory, methods and Applications*, 22 (4), Pp. 499-503.
- LaSalle, J., 1976. The stability of dynamical systems, in *Proceedings of the*

- CBMS-NSF Regional Conference Series in Applied Mathematics 25, SIAM, Philadelphia, Pa, USA.
- Martcheva, M., 2015. An Introduction to Mathematical Epidemiology. Texts in Applied Mathematics, 61, Springer, New York, NY, USA.
- May, R.C., 1973. Stable limit cycles in pre-predator population. *A comment*, Science, Pp. 1074.
- Pielou, E.C., 1969. An introduction to Mathematical Ecology. Wiley-InterScience, Hoboken, NJ, USA.
- Rabies fact sheet, 2016. <http://www.who.int/mediacentre/factsheets/fs099/en/>.
- Rupprecht, C.E., Briggs, D., Brown, C.M., Franka, R., Katz, S.L, Kerr, H.D., 2010. Use of A Reduced (4-Dose) Vaccine Schedule for Postexposure Prophylaxis to Prevent Human Rabies: Recommendations of The Advisory Committee on Immunization Practices, Department of Health and Human Services, Centers for Disease Control and Prevention.
- Samia, B., Sajjad, A., Kamal, S., Muhammad, A., 2018. Exact solution to nonlinear biological population model with fractional order. *Thermal Science Journal*, Special Issue, 10 pages
- Shah, K., Ali, A., Bushnaq, S., 2018. Hyers–Ulam stability analysis to implicit Cauchy problem of fractional differential equations with impulsive conditions. *Math. Methods Appl. Sci.*, 41(17), 8329–8343.
- Sharomi, O., Malik, T., 2017. Optimal control in epidemiology. *Annals of Operations Research*, 251 (1-2), Pp. 5571.
- Simon A. Levin, Thomas G. Hallam and Louis J. Gross (Editors). 1989. *Applied Mathematical Ecology*. *Biomathematics*, 18, Pp. 491.
- Smith, G.C., Cheeseman, C.L., 2002. A mathematical model for the control of diseases in wildlife populations: culling, vaccination and fertility control. *Ecological Modelling*, 150 (1-2), Pp. 4553.
- Tchuenche, J.M., Bauch, C.T., 2012. Can culling to prevent monkeypox infection be counter-productive? Scenarios from a theoretical model. *Journal of Biological Systems*, 20 (3), Pp. 259-283.
- Wang, X., Lou, J., 2008. Two dynamic models about rabies between dogs and human. *Journal of Biological Systems*, 16 (4), Pp. 519-529.
- Wiraningsih, E.D., Agosto, F., Aryati, L., 2015. Stability analysis of rabies model with vaccination effect and culling in dogs. *Applied Mathematical Sciences*, 9 (77-80), Pp. 3805-3817.
- Yang, W., Lou, J., 2009. The dynamics of an interactional model of rabies transmitted between human and dogs. *Bollettino della Unione Matematica Italiana, Serie 9*, 2 (3), Pp. 591-605.
- Yusuf, T.T., Benyah, F., 2012. Optimal control of vaccination and treatment for an SIR epidemiological model. *World Journal of Modelling and Simulation*, 8 (3), Pp. 194-204.
- Zhang, J., Jin, Z., Sun, G.Q., Zhou, T., Ruan, S., 2011. Analysis of rabies in China: transmission dynamics and control. *PLoS ONE*, 6 (7), Pp. e20891.
- Zhang, T., Wang, K., Zhang, X., Jin, Z., 2015. Modeling and analyzing the transmission dynamics of HBV epidemic in Xinjiang, China. *PLoS ONE*, 10 (9), Article ID e138765.
- Zinsstag, J., Durr, S., Penny, M.A., 2009. Transmission dynamics and economics of rabies control in dogs and humans in an African city. *Proceedings of the National Academy of Sciences of the United States of America*, 106 (35), Pp. 14996-15001.

