



Print ISSN : 2521-0831
Online ISSN : 2521-084X

CODEN: MSMADH

SOME CONTRIBUTION OF SOFT PRE-OPEN SETS TO SOFT W-HAUSDORFF SPACE IN SOFT TOPOLOGICAL SPACES

Arif Mehmood Khattak^{1*}, Ishfaq Ahmed², Zaheer Anjum³, Muhammad Zamir³, Fahad Jamal²


¹Riphah International University Islamabad Math and Stastics department (Pakistan)

²Qurtuba University of Science and Information Technology Peshawar Math department (Pakistan)

³Department of Mathematics Bannu University of Sciences and Technology (Pakistan)

*Corresponding Author's E-mail: mehdaniyal@gmail.com

This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited



ARTICLE DETAILS

ABSTRACT

Article History:

Received 12 November 2017
Accepted 12 December 2017
Available online 1 January 2018

In this article the notion of soft W-Hausdorff or soft $W-T_2$ structure in soft topological spaces is proclaimed with respect to soft pre-open sets while using the ordinary points of Soft Topology. That's is why it is named as soft P-W-Hausdorff or soft P-W- T_2 structure. Some sub-spaces of soft P-W- T_2 structure is also reflected. Product of these spaces is also avail.

KEYWORDS

Land use/Land cover Change, Wetland, Remote Sensing, Landsat TM/ETM+, Marko.

1. INTRODUCTION

Based on a study, general topology plays an attractive role in space time geometry as well as different branches of pure and applied mathematics [1]. In Separation Axioms we discuss points of the space [2]. It shows the points are separated by neighbour-hood [3]. Based on a research, when we are interested to know the distance among the points that are separated from each other, then in that case the concept of separation axioms will come in play [4-6]. A scholar said that most of the real-life problem have various uncertainties [7]. A number of theory have been proposed for handling with uncertainties in an efficient way in 1999 ,where a researcher was first who given birth to a novel concept of soft set theory which is completely a new approach for modelling vagueness and uncertainty in 2011, another researcher define soft topological spaces and studied separation axioms and section two of this paper preliminary definition regarding soft sets and soft topological spaces are given in section three of this paper the concept of soft W-Housdroffness in soft topological spaces is introduced by referring the definition of soft P-open sets [8-10]. Throughout this paper \tilde{X} denotes the father set and E denotes the set of parameters for the father \tilde{X} .

2. PRELIMINARIES

Definition 1: Let \tilde{X} be the father set and \tilde{E} be a set of parameters. Let $P(\tilde{X})$ denotes the power set of and \tilde{A} be a nonempty subset of \tilde{E} . A pair (F, \tilde{A}) denoted by $F_{\tilde{A}}$ Is called a soft set over \tilde{X} where F is a mapping given by $F: \tilde{A} \rightarrow P(\tilde{X})$. In other words a soft set over \tilde{X} is parameterized family of subsets of the universe \tilde{X} for a particular $e \in \tilde{A}$, $F(e)$ may be considered the set of e-approximate elements of the soft set equation (F, \tilde{A}) if $e \notin \tilde{A}$

Then $F(e) = \tilde{\emptyset}$ $F_{\tilde{A}} = \{F(e); e \in \tilde{A} \subseteq \tilde{E}; F: \tilde{A} \rightarrow P(\tilde{X})\}$

The family of all these soft set over \tilde{X} w.r.t the parameter set \tilde{E} is denoted $SS(\tilde{X})_{\tilde{E}}$

Definition 2: Let $F_{\tilde{A}}, G_{\tilde{B}} \in SS(\tilde{X})_{\tilde{E}}$. then $F_{\tilde{A}}$ is soft sub set of $G_{\tilde{B}}$ is denoted by $F_{\tilde{A}} \subseteq G_{\tilde{B}}$ if

- (1) $\tilde{A} \subseteq \tilde{B}$ and
- (2) $F(e) \subseteq G(e), \forall e \in \tilde{A}$

In this case $F_{\tilde{A}}$ is said to be a soft subset of $G_{\tilde{B}}$ and $G_{\tilde{B}}$ is said to be soft super set of $F_{\tilde{A}}$, $G_{\tilde{B}} \supseteq F_{\tilde{A}}$

Definition 3: Two soft subset $F_{\tilde{A}}$ and $G_{\tilde{B}}$ over a common father set \tilde{X} are said to be equal if $F_{\tilde{A}}$ is soft subset of $G_{\tilde{B}}$ and $G_{\tilde{B}}$ is a soft subset of $F_{\tilde{A}}$.

Definition 4: Let τ be the collection of soft sets over \tilde{X} , then τ is said to be a soft topology on \tilde{X} , if

1. \emptyset, \tilde{X} belong to τ
 2. The union of any number of soft sets in τ belongs to τ
 3. The intersection of any two soft sets in τ belong to τ
- The triplet (\tilde{X}, F, E) is called a soft topological space.

Definition 5: Let (F, A) be any soft set of a soft topological space (\tilde{X}, τ, A) then (F, A) is called

- i. 1) (F, A) soft pre-open set of \tilde{X} if $(F, A) \subseteq \text{int}(\text{cl}((F, A)))$ and
 - ii. 2) (F, A) soft pre-closed set of \tilde{X} if $(F, A) \supseteq \text{cl}(\text{int}((F, A)))$
- The set of all pre-open soft sets is denoted by $POS(\tilde{X})$

Definition 6: Complement of a soft (F, \tilde{A}) set denoted by $(F, \tilde{A})^c$ denoted by $(F, \tilde{A})^c = \{F^c(e); \tilde{A} \rightarrow P(\tilde{X})\}$ is a mapping given by $F^c(e) = \tilde{X} - F(e); \forall e \in \tilde{A}$ and F^c is called the soft complement function of F . clearly $(F^c)^c = F$. clearly $(F^c)^c = F$.

Definition 7: A soft set (F, \tilde{A}) over \tilde{X} is said to be a null soft set denoted by $\tilde{\emptyset}$ or $\tilde{\emptyset}_{\tilde{A}}$ if $\forall e \in \tilde{A}, F(e) = \emptyset$

Definition 8:[6] A soft set (F, \tilde{A}) over \tilde{X} is said in absolute soft set denoted by \tilde{A} or $X_{\tilde{A}}$ IF $\forall e \in \tilde{A}, F(e) = \tilde{X}$ clearly we have $\tilde{X}_{\tilde{A}}^c = \tilde{\emptyset}_{\tilde{A}}$ and $\tilde{\emptyset}_{\tilde{A}}^c = X_{\tilde{A}}$

Definition 9: The union of two soft set (F, \tilde{A}) and (G, \tilde{B}) over the common universe \tilde{X} is the soft set (H, \tilde{C}) where $\tilde{C} = \tilde{A} \cup \tilde{B}$ and $\forall e \in \tilde{C}$

$$H(e) = \begin{cases} F(e), e \in \tilde{A} - \tilde{B} \\ G(e), e \in \tilde{B} - \tilde{A} \\ F(e) \cup G(e), e \in \tilde{A} \cap \tilde{B} \end{cases}$$

Definition 10: The intersection of two soft set (F, \tilde{A}) and (G, \tilde{B}) over the common universe \tilde{X} is the soft set (H, \tilde{C}) where $\tilde{C} = \tilde{A} \cap \tilde{B}$ and $\forall e \in \tilde{C}$
 $\tilde{H}(e) = F(e) \cap G(e)$.

Definition 11: Let $(\tilde{X}, \tau, \tilde{E})$ be soft topological space $(F, E) \in SS(X)_E$ and Y be a non null subset of X then the soft subset of (F, E) over Y denoted by (F_Y, \tilde{E}) is defined as follows $F_Y(e) = \tilde{Y} \cap F(e) \forall e \in \tilde{E}$ In other words $(F_Y, \tilde{E}) = \tilde{Y}_E \cap (F, \tilde{E})$

Definition 12: Let $(\tilde{X}, \tau, \tilde{E})$ be soft topological space (F, \tilde{E}) and \tilde{Y} be a non-null subset of \tilde{X} . then $\{(F, \tilde{E}) : (F, \tilde{E}) \in \tau\}$ is said to be the relative soft topology on Y and $(\tilde{Y}, \tau_Y, \tilde{E})$ is called a soft subspace of $(\tilde{X}, \tau, \tilde{E})$

Definition 13: $F_A \in SS(X)_E$ and $G_B \in SS(Y)_k$. The Cartesian product $(F_A \odot G_B)(e, k) = (F_A(e) \times G_B(k)) \forall (e, k) \in \tilde{A} \times \tilde{B}$. According to this definition $(F_A \odot G_B)$ is soft set over $\tilde{X} \times \tilde{Y}$ and its parameter set is $\tilde{E} \times \tilde{K}$

Definition 14: $(\tilde{X}, \tau_X, \tilde{E})$ and $(\tilde{Y}, \tau_Y, \tilde{K})$ BE two soft topological spaces the soft product topology $\tau_X \odot \tau_Y$ over $\tilde{X} \times \tilde{Y}$ w.r.t $\tilde{E} \times \tilde{K}$ is the soft topology having the collection $\{F_E \odot G_K / F_A \in \tau_X, G_K \in \tau_Y\}$ is the basis.

3. SOFT P-W-HAUSDORFF SPACES

This section is dedicated to soft Pre-W-T₂Space. Different results are deeply discussed related to this soft topological structure with the application of soft pre-open sets while using ordinary points of Soft Topology.

Definition 15: A soft topological space $(\tilde{X}, \tau, \tilde{E})$ is said to be soft pre W-Hausdorff space of type 1 denoted by $(PSW - H)_1$ if for every $e_1, e_2 \in \tilde{E}, e_1 \neq e_2$ there exist soft pre-open sets $(F, \tilde{A}), (G, \tilde{B})$ such that $F_A(e_1) = X, G_B(e_2) = \tilde{X}$ and $(F, \tilde{A}) \cap (G, \tilde{B}) = \tilde{\phi}$.

Proposition 1. Soft subspace of a $(PSW - \tilde{H})_1$ space is soft $(PSW - H)_1$

Proof: let $(\tilde{X}, \tau, \tilde{E})$ be a $(PSW - \tilde{H})_1$ space. Let \tilde{Y} be a non-vacuous subset of \tilde{X} . Let $(\tilde{Y}, \tau_Y, \tilde{E})$ be soft subspace of $(\tilde{X}, \tau, \tilde{E})$ where $\{\tau_Y = F_Y, \tilde{E}\} : (F, \tilde{E}) \in \tau$ is the relative soft topology on \tilde{Y} . Considered $e_1, e_2 \in \tilde{E}, e_1 \neq e_2$, there exist soft Pre-open sets $(F, \tilde{A}), (G, \tilde{B})$ such that $F_A(e_1) = \tilde{X}, G_B(e_2) = \tilde{X}$ and $F_A \cap G_B = \tilde{\phi}$

Therefore, $((F_A)_Y, E), ((G_B)_Y, \tilde{E}) \in \tau_Y$
 $= \tilde{Y} \cap \tilde{X}$
 $= \tilde{Y}$

$((G_B)_Y, (e_2) = \tilde{Y} \cap G_B(e_2)$
 $= \tilde{Y} \cap \tilde{X}$
 $= \tilde{Y}$

$(F_A)_Y \cap (G_B)_Y(e) = (F_A \cap G_B)_Y(e)$
 $= \tilde{Y} \cap (F_A \cap G_B)_Y(e)$
 $= \tilde{Y} \cap \tilde{\phi}(e)$
 $= \tilde{Y} \cap \tilde{\phi}$
 $= \tilde{\phi}$

$(F_A)_Y \cap (G_B)_Y = \tilde{\phi}$
Hence (Y, τ_Y, E) is soft pre $(SW - \tilde{H})_1$ that is $(PSW - \tilde{H})_1$

Definition 16: Let $(\tilde{X}, \tau, \tilde{E})$ be a soft topological and $\tilde{H} \subseteq \tilde{E}$ then let $(\tilde{X}, \tau_{\tilde{H}}, \tilde{E})$ is called soft subspace of $(\tilde{X}, \tau, \tilde{E})$ relative to the parameter set H where $\tau_{\tilde{H}} = \{(F_A / \tilde{H} : \tilde{H} \subseteq \tilde{A} \subseteq \tilde{E}; (F, \tilde{A}) \text{ is soft Pre - open set}\}$ and $= \{(F_A) / H\}$ is the restriction map on \tilde{H}

Proposition 2. Soft p -subspace of a $(PSW - H)_1$ space is $(PSW - H)_1$

Proof: let $(\tilde{X}, \tau, \tilde{E})$ be a $(PSW - H)_1$ space. Let (Y, τ_H, \tilde{E}) be soft p -subspace of $(\tilde{X}, \tau, \tilde{E})$ relative parameter set H where $H = \{(F_A / \tilde{H} : \tilde{H} \subseteq \tilde{A} \subseteq \tilde{E}; \text{where } (F, \tilde{A}) \text{ is soft Pre - open set } \in \tau\}$ Consider $h, h \in \tilde{H} \quad h_1 \neq h_2$ then $h, h \in \tilde{E}$ Therefore, there exists Pre-open sets $(F, \tilde{A}), (G, \tilde{B})$ such that $F_A(h_1) = X, G_B(h_2) = X$ and $(F, \tilde{A}) \cap (G, \tilde{B}) = \tilde{\phi}$

Therefore $(F_A) / H, (G_B) / \tilde{H} \in \tau_H$
 $(F_A) / \tilde{H}(h_1) = F_A(h_1) = \tilde{X}$

$(G_B) / \tilde{H}(h_2) = G_B(h_2) = \tilde{X}$
 $(F_A) / \tilde{H} \cap (G_B) / \tilde{H} = (F_A \cap G_B) / \tilde{H}$
 $= \tilde{\phi} / H$
 $= \tilde{\phi}$

Hence (\tilde{X}, τ_H, H) is $(SSW - H)_1$

Proposition 3. Product of two $(PSW - H)_1$ spaces $(PSW - H)_1$

Proof: let $(\tilde{X}, \tau_X, \tilde{E})$ and let $(\tilde{Y}, \tau_Y, \tilde{K})$ be two $(PSW - H)_1$ spaces Consider two distinct points (e_1, \tilde{K}_1) and $(e_2, \tilde{K}_2) \in \tilde{E} \times \tilde{K}$

Either $e_1 \neq e_2$ or $\tilde{K}_1 \neq \tilde{K}_2$ Assume $e_1 \neq e_2$ $(\tilde{X}, \tau_X, \tilde{E})$ is $(PSW - H)_1$ there exist soft Pre-open sets $(F, \tilde{A}), (G, \tilde{B})$ such that $F_A(e_1) = \tilde{X}, G_B(e_2) = \tilde{X}$ and $(F, \tilde{A}) \cap (G, \tilde{B}) = \tilde{\phi}$ Therefore $F_A \odot Y_K, G_B \odot \tilde{Y}_K \in \tau_X \odot \tau_Y$
 $(F_A \odot Y_K)(e_1, \tilde{K}_1) = F_A(e_1) \times \tilde{Y}_K(\tilde{K}_1)$
 $= \tilde{X} \times \tilde{Y}$

$(G_B \odot \tilde{Y}_K)(e_2, \tilde{K}_2) = G_B(e_2) \times \tilde{A}_K(\tilde{K}_2)$
 $= \tilde{X} \times \tilde{Y}$

If for any

$(e, k) \in \tilde{E} \times \tilde{K}, (F_A \odot Y_K)(e, k) \neq \tilde{\phi}$
 $\Rightarrow F_A(e) \times \tilde{Y}_K(k) \neq \tilde{\phi}$
 $\Rightarrow F_A(e) \times \tilde{Y}_K \neq \tilde{\phi}$
 $\Rightarrow F_A(e) \neq \tilde{\phi}$
 $\Rightarrow G_B(e) = \tilde{\phi}$ Since $F_A \cap G_B = \tilde{\phi}$
 $\Rightarrow F_A(e) \cap G_B(e) = \tilde{\phi}$
 $\Rightarrow F_A(e) \times \tilde{Y}_K(k) = \tilde{\phi}$

$(G_B \odot \tilde{Y}_K)(e, k) = \tilde{\phi}$

$(F_A \odot \tilde{Y}_K) \cap (G_B \odot \tilde{Y}_K) = \tilde{\phi}$
Assume $\tilde{K}_1 \neq \tilde{K}_2$ $(\tilde{Y}, \tau_Y, \tilde{K})$ is $(PSW - H)_1$ there exist Pre-open sets $(F, \tilde{A}), (G, \tilde{B})$ $F_A(K_1) = Y, G_B(K_2) = \tilde{Y}$ and $(F, \tilde{A}) \cap (G, \tilde{B}) = \tilde{\phi}$

$(\tilde{X}_E \odot F_A)(e_1, K_1) = \tilde{X}_E(e_1) \times F_A(K_1)$
 $= \tilde{X} \times \tilde{Y}$

$(\tilde{X}_E \odot G_B)(e_2, \tilde{K}_2) = (\tilde{X}_E(e_2) \times G_B(K_2))$
 $= \tilde{X} \times \tilde{Y}$

If for any $(e, k) \in \tilde{E} \times \tilde{K}$

$\Rightarrow (\tilde{X}_E(e) \times F_A(k)) \neq \tilde{\phi}$
 $\Rightarrow \tilde{X} \times F_A(k) \neq \tilde{\phi}$
 $\Rightarrow F_A(k) \neq \tilde{\phi}$

$G_B(k) = \tilde{\phi}$ Since $(F, \tilde{A}), (G, \tilde{B}) = \tilde{\phi}$ that is $F_A \cap G_B = \tilde{\phi}$

$\Rightarrow F_A(k) \cap G_B(K) = \tilde{\phi}$
 $\Rightarrow \tilde{X}_E(e) \times G_B(e) = \tilde{\phi}$
 $\Rightarrow (\tilde{X}_E \odot G_B)(e, k) = \tilde{\phi}$

$(\tilde{X}_E \odot F_A) \cap (\tilde{X}_E \odot G_B) = \tilde{\phi}$

Hence $\tilde{X} \times \tilde{Y} (\tau_X \odot \tau_Y, \tilde{E} \times \tilde{K})$ is $(PSW - H)_1$

Definition 17: A soft topological space (X, τ, E) is said to be soft Semi W-hausdorff that is P-W-Hausdorff space of type 2 denoted by $(PSW - H)_2$

If for every $e_1, e_2 \in \tilde{E}, e_1 \neq e_2$ there exists soft Per-open sets $(F, \tilde{E}), (G, \tilde{E})$ such that

$F_e(e_1) = \tilde{X}, G_e(e_2) = \tilde{X}$ and $F_e \cap G_e = \tilde{\phi}$

Proposition 4. Soft subspace of a $(PSW - \tilde{H})_2$ space is $(PSW - \tilde{H})_2$

Proof: Let $(\tilde{X}, \tau, \tilde{E})$ be a $(PSW - \tilde{H})_2$ space. Let \tilde{Y} be a Non-null subset of \tilde{X} . Let $(\tilde{Y}, \tau_Y, \tilde{E})$ be a soft subspace Of $(\tilde{X}, \tau, \tilde{E})$ where $\tau_Y = \{(F_Y, \tilde{E}) : \text{where } (F, \tilde{E}) \text{ is soft Pre - open sets } \in \tau\}$ is the relative soft topology on \tilde{Y} . Considered $e_1, e_2 \in \tau, e_1, e_2 \neq \tau$ There exist soft Pre-open sets $(F, \tilde{E}), (G, \tilde{E})$ such that $F_e(e_1) = \tilde{X}, G_e(e_2) = \tilde{X}$ and $(F, \tilde{E}), (G, \tilde{E}) = \tilde{\phi}$ that is $F_e \cap G_e = \tilde{\phi}$.

Therefore $((F_e)_Y, \tilde{E}), ((G_e)_Y, \tilde{E}) \in \tau_Y$

Also $(F_e)_Y(e_1) = \tilde{Y} \cap F_e(e_1)$
 $= \tilde{Y} \cap \tilde{X}$
 $= \tilde{Y}$

$(G_e)_Y(e_2) = \tilde{Y} \cap G_e(e_2)$
 $= \tilde{Y} \cap \tilde{X}$
 $= \tilde{Y}$

$((F_e)_Y \cap (G_e)_Y)(e) = ((F_e \cap G_e)_Y)(e)$
 $= \tilde{Y} \cap (F_e \cap G_e)_Y(e)$
 $= \tilde{Y} \cap \tilde{\phi}$

$$\begin{aligned}
 &= \tilde{Y} \cap \tilde{\emptyset} \\
 &= \tilde{\emptyset} \\
 (F_{\tilde{A}})_{\tilde{Y}} \cap (G_{\tilde{B}})_{\tilde{Y}} &= \tilde{\emptyset} \\
 \text{Hence } (\tilde{Y}, \tau_{\tilde{Y}}, \tilde{E}) &\text{ is } (SSW - \tilde{H})_2
 \end{aligned}$$

Proposition 5. Soft p-subspace of a $(PSW - \tilde{H})_2$ is $(PSW - \tilde{H})_2$

Proof: Let $(\tilde{X}, \tau, \tilde{E})$ be a $(PSW - \tilde{H})_2$ space. Let $\tilde{H} \subseteq \tilde{E}$
 Let $(\tilde{X}, \tau_{\tilde{H}}, \tilde{H})$ be a soft p-subspace of $(\tilde{X}, \tau, \tilde{E})$, relative to the parameter set \tilde{H} where $\{(F_{\tilde{A}})/\tilde{H}: \tilde{H} \subseteq \tilde{A} \subseteq \tilde{E}, (F, \tilde{A}) \in \tau\}$ where (F, \tilde{A}) is soft Pre – open set

Consider $h_1, h_2 \in \tilde{H}, h_1 \neq h_2$.
 Then $h_1, h_2 \in \tilde{H}$ there exists soft pre-open set $(F, \tilde{E}), (G, \tilde{E})$
 Such that $F_{\tilde{E}}(h_1) = \tilde{X}, G_{\tilde{E}}(h_2) = \tilde{X}$ and $(F, \tilde{E}), (G, \tilde{E}) = \tilde{\emptyset}$ that is $F_{\tilde{E}} \cap G_{\tilde{E}} = \tilde{\emptyset}$
 Therefore $(F_{\tilde{A}})/\tilde{H}, ((G_{\tilde{B}})/\tilde{H}) \in \tau_{\tilde{H}}$
 Also $((F_{\tilde{E}})/\tilde{H})(h_1) = F_{\tilde{E}}(h_1) = \tilde{X}$
 $((G_{\tilde{E}})/\tilde{H})(h_2) = G_{\tilde{E}}(h_2) = \tilde{X}$ &
 $((F_{\tilde{E}})/\tilde{H}) \cap ((G_{\tilde{E}})/\tilde{H}) = (F_{\tilde{E}} \cap G_{\tilde{E}})/\tilde{H}$
 $= \tilde{\emptyset}/\tilde{H}$
 $= \tilde{\emptyset}$

Hence $(\tilde{X}, \tau, \tilde{E})$ is $(SSW - H)_2$

Proposition 6. Product of two $(PSW - H)_2$ spaces is $(PSW - H)_2$

Proof: Let $(\tilde{X}, \tau_{\tilde{X}}, \tilde{E})$ & $(\tilde{Y}, \tau_{\tilde{Y}}, \tilde{K})$ be two $SSW - H)_2$ Soft spaces. Consider two distinct points $(e_1, k_1), (e_2, k_2) \in \tilde{E} \times \tilde{K}$
 Either $e_1 \neq e_2$ or $k_1 \neq k_2$
 Assume $e_1 \neq e_2$ Since $(\tilde{X}, \tau_{\tilde{X}}, \tilde{E})$ is $(PSW - H)_2$, there Soft Pre-open sets $(F, \tilde{E}), (G, \tilde{E}) \in \tau_{\tilde{X}}$ such that $F_{\tilde{E}}(e_1) = \tilde{X}, G_{\tilde{E}}(e_2) = \tilde{X}$ and $(F, \tilde{E}), (G, \tilde{E}) = \tilde{\emptyset}$ that is $F_{\tilde{E}} \cap G_{\tilde{E}} = \tilde{\emptyset}$
 Therefore $F_{\tilde{E}} \odot \tilde{Y}_{\tilde{K}}, G_{\tilde{E}} \odot \tilde{Y}_{\tilde{K}} \in \tau_{\tilde{X}} \odot \tau_{\tilde{Y}}$
 $(F_{\tilde{E}} \odot \tilde{Y}_{\tilde{K}})(e_1, k_1) = F_{\tilde{E}}(e_1) \times \tilde{Y}_{\tilde{K}}(k_1)$
 $= \tilde{X} \times \tilde{Y}$
 $(G_{\tilde{E}} \odot \tilde{Y}_{\tilde{K}})(e_2, k_2) = G_{\tilde{E}}(e_2) \times \tilde{Y}_{\tilde{K}}(k_2)$
 $= \tilde{X} \times \tilde{Y}$
 If for any $(e, k) \in (\tilde{E} \times \tilde{K}), (F_{\tilde{E}} \odot \tilde{Y}_{\tilde{K}})(e, k) \neq \tilde{\emptyset}$
 $\Rightarrow F_{\tilde{E}}(e) \times \tilde{Y}_{\tilde{K}}(k) \neq \tilde{\emptyset}$
 $\Rightarrow F_{\tilde{E}}(e) \times \tilde{Y} \neq \tilde{\emptyset}$
 $\Rightarrow F_{\tilde{E}}(e) \neq \tilde{\emptyset}$
 $\Rightarrow G_{\tilde{E}}(e) = \tilde{\emptyset}$ (since $F_{\tilde{E}} \cap G_{\tilde{E}} = \tilde{\emptyset} \Rightarrow F_{\tilde{E}}(e) \cap G_{\tilde{E}}(e) = \tilde{\emptyset}$)
 $\Rightarrow G_{\tilde{E}}(e) \times \tilde{Y}_{\tilde{K}}(k) = \tilde{\emptyset}$
 $\Rightarrow G_{\tilde{E}} \odot \tilde{Y}_{\tilde{K}}(e, k) = \tilde{\emptyset}$
 $\Rightarrow (F_{\tilde{E}} \odot \tilde{Y}_{\tilde{K}}) \cap (G_{\tilde{E}} \odot \tilde{Y}_{\tilde{K}}) = \tilde{\emptyset}$
 In a similar fashion, one can prove the case when $k_1 \neq k_2$
 Hence $(\tilde{X} \times \tilde{Y}, \tau_{\tilde{X}} \odot \tau_{\tilde{Y}}, \tilde{E} \times \tilde{K})$ is $(PSW - H)_2$.

*

4. CONCLUSION

In this article the notion of Soft Pre-W-Hausdorff spaces is presented with respect to ordinary points of the soft topological spaces and some basic properties regarding this concept are established in a better way. I have fastidiously studied numerous homes on the behalf of Soft Topology. And lastly, I determined that soft Topology is totally linked or in other sense we can correctly say that Soft Topology (Separation Axioms) are connected with structure. Provided if it is related with structures then it gives the idea of non-linearity beautifully. In other ways we can rightly say Soft Topology is somewhat directly proportional to non-linearity. Although we use non-linearity in Applied Math. So, it is not wrong to say that Soft Topology is applied Math in itself. It means that Soft Topology has the taste of both of pure and applied math. In future I will discuss Separation Axioms in Soft Topology With respect to soft points.

REFERENCES

[1] Ali, M.I., Feng, F., Liu, X., Min, W.K., Shabir, M. 2009. On some new operations in soft set theory. Computers and Mathematics with Applications, 57, 1547-1553.

[2] Babitha, K.V., Sunil, J.J. 2010. Soft set relations and functions, Computers and Mathematics with Applications, 60, 1840-1848.
 [3] Chang, C.L. 1968. Fuzzy Topological Spaces, Journal of Mathematical Analysis and Applications, 24, (1): 182-190.
 [4] Gantner, T.E., Steinlage R.C., Warren, R.H. 1978. Compactness in Fuzzy Topological Spaces, Journal of Mathematical Analysis and Applications, 62, 547-562.
 [5] Maji, P.K., Biswas, R., Roy, A.R. 2003. Soft Set Theory, Computers and Mathematics with Applications, 45, (4-5): 555-562.
 [6] Molodstov, D. 1999. Soft Set Theory – First Results, Computers and Mathematics with Applications, 37, (4-5): 19-31.
 [7] Shabir, M., Naz, M. 2011. On Soft Topological Spaces, Computers and Mathematics with Applications, 61, (7): 1786-1799.
 [8] Zou, Z., Xiao, Z. 2008. Data analysis approach of soft sets under incomplete information, Knowledge-Based System, 21, 941-945.
 [9] Hussain, S., Ahmad, B. 2015. Soft separation axioms in soft topological space submitted to elsvier.
 [10] Arockiarani, Lancy, A.A. 2013. Generalized soft gβ-closed sets and soft gβ-closed sets in soft topological spaces, IJMRS, 4 (2): 1-7.

